

NATIONAL DRUG CODE (NDC) AND BILLING AND DIAGNOSIS CODES FOR KEYTRUDA

The following codes may be relevant when billing for KEYTRUDA and its administration. This information is current as of February 2021. The information provided here is compiled from sources believed to be accurate, but Merck makes no representation that it is accurate. Information about HCPCS codes is based on guidance issued by the CMS applicable to Medicare Part B and may not apply to other public or private payers. Consult the relevant manual and/or other guidelines for a description of each code to determine the appropriateness of its use and for information on additional codes. Diagnosis codes should be selected only by a health care professional. This information is subject to change. Merck cautions that payer-coding requirements vary and can frequently change, so it is important to regularly check with each payer or, where applicable, the Medicare Administrative Contractor as to payer-specific requirements.

You are solely responsible for determining the appropriate codes and for any action you take in billing. The information provided here is not intended to be definitive or exhaustive, and is not intended to replace the guidance of a qualified professional advisor. Diagnosis codes should be selected only by a health care professional. Merck and its agents make no warranties or guarantees, expressed or implied, concerning the accuracy or appropriateness of this information for your particular use given the frequent changes in public and private payer billing. The use of this information does not guarantee payment or that any payment received will cover your costs.

NDC and Packaging Information

The NDC is typically required when submitting a claim with a miscellaneous HCPCS code. Please consult with the payer to understand specific billing requirements.

PRODUCT	
KEYTRUDA [®] (pembrolizumab) injection 100 mg	
PACKAGE	NDC
Carton containing one 100 mg/4 mL (25 mg/mL), single-dose vial	0006-3026-02
Carton containing two 100 mg/4 mL (25 mg/mL), single-dose vials	0006-3026-04

Please note: The NDCs above are the billable NDCs that appear on the cartons. The NDC on the vial should not be used for billing purposes.

Billing Codes

Below is a list of possible codes that could be relevant for KEYTRUDA and its administration. Please consult with the applicable payer to understand the payer's specific billing requirements.

HCPCS Code for KEYTRUDA¹

HCPCS CODE	DESCRIPTOR
J9271	Injection, pembrolizumab, 1 mg

Revenue Code—For Use in the Hospital Setting²

REVENUE CODE	DESCRIPTOR
0636	Drug requiring detailed coding

Current Procedural Terminology (CPT[®])^a Code for Administration^{3,4}

CPT CODE	DESCRIPTOR
96413 (Some payers may utilize 96365. Check with the applicable payer.)	Injection and Intravenous Infusion Chemotherapy and Other Highly Complex Drug or Highly Complex Biologic Agent Administration

^aCPT Copyright 2019 American Medical Association. All rights reserved.

Before prescribing KEYTRUDA, please read the Selected Safety Information on pages 5-8 and the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA FOR RECURRENT OR METASTATIC CUTANEOUS SQUAMOUS CELL CARCINOMA (cSCC)

The following codes as of February 2021 are provided as a reference and may be relevant when billing for KEYTRUDA and its administration. Consult the relevant manual and/or other guidelines for a description of each code to determine the appropriateness of its use and for information on additional codes. Diagnosis codes should be selected only by a health care professional. You are solely responsible for determining the appropriate codes and for any action you take in billing.

When submitting a claim for KEYTRUDA, always verify coding requirements with the relevant payer. Coding requirements may vary by insurer or plan; please refer to the payer-specific policies to understand what may be covered.

Check with the relevant payer regarding guidance on which diagnoses they will recognize and the applicability of secondary codes. Health care professionals are solely responsible for selecting codes that appropriately reflect the patient's diagnosis, the services rendered, and the applicable payers' guidelines.

Providers should document the diagnosis with a sufficiently high degree of specificity based on the information available to enable the identification of the most appropriate code. Although CMS has said that an unspecified code may be appropriate in some cases, CMS has advised that you should always code with as much specificity as possible consistent with the clinical documentation.

Merck and its agents make no warranties concerning the accuracy or appropriateness of this information for your particular use given the frequent changes in public and private payer billing. Merck cautions that payer-coding requirements vary and can frequently change, so it is important to regularly check with each payer or, where applicable, the Medicare Administrative Contractor as to payer-specific requirements. The use of this information does not guarantee payment or that any payment received will cover your costs.

Indication

KEYTRUDA is indicated for the treatment of patients with recurrent or metastatic cutaneous squamous cell carcinoma (cSCC) that is not curable by surgery or radiation.

FDA-Approved Dosing

The FDA-approved dose of KEYTRUDA is either 200 mg administered as an intravenous infusion over 30 minutes every 3 weeks or 400 mg administered as an intravenous infusion over 30 minutes every 6 weeks, until disease progression, unacceptable toxicity, or up to 24 months.

KEYTRUDA is indicated for use at an additional recommended dosage of 400 mg every 6 weeks for all approved adult indications. This indication is approved under accelerated approval based on pharmacokinetic data, the relationship of exposure to efficacy, and the relationship of exposure to safety. Continued approval for this dosing may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Possible relevant diagnosis codes for cSCC

Learn more about each diagnosis code and descriptor by selecting from the series options below.

C44: Other and Unspecified Malignant Neoplasm of Skin⁴

The C44 series:

- *Excludes:* Kaposi's sarcoma of skin, malignant melanoma of skin, malignant neoplasm of skin of genital organs, Merkel cell carcinoma

ICD-10-CM CODE	DESCRIPTOR
C44.0	Other and unspecified malignant neoplasm of skin of lip

Before prescribing KEYTRUDA, please read the Selected Safety Information on pages 5-8 and the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg

KEYTRUDA FOR RECURRENT OR METASTATIC CUTANEOUS SQUAMOUS CELL CARCINOMA (cSCC) (continued)

C44.12: Squamous Cell Carcinoma of Skin of Eyelid, Including Canthus⁴

ICD-10-CM CODE	DESCRIPTOR
C44.121	Squamous cell carcinoma of skin of unspecified eyelid, including canthus
C44.1221	Squamous cell carcinoma of skin of right upper eyelid, including canthus
C44.1222	Squamous cell carcinoma of skin of right lower eyelid, including canthus

C44.129: Squamous Cell Carcinoma of Skin of Left Eyelid, Including Canthus⁴

ICD-10-CM CODE	DESCRIPTOR
C44.1291	Squamous cell carcinoma of skin of left upper eyelid, including canthus
C44.1292	Squamous cell carcinoma of skin of left lower eyelid, including canthus

C44.22: Squamous Cell Carcinoma of Skin of Ear and External Auricular Canal⁴

ICD-10-CM CODE	DESCRIPTOR
C44.221	Squamous cell carcinoma of skin of unspecified ear and external auricular canal
C44.222	Squamous cell carcinoma of skin of right ear and external auricular canal
C44.229	Squamous cell carcinoma of skin of left ear and external auricular canal

C44.32: Squamous Cell Carcinoma of Skin of Other and Unspecified Parts of Face⁴

ICD-10-CM CODE	DESCRIPTOR
C44.320	Squamous cell carcinoma of skin of unspecified parts of face
C44.321	Squamous cell carcinoma of skin of nose
C44.329	Squamous cell carcinoma of skin of other parts of face

C44.4: Other and Unspecified Malignant Neoplasm of Skin of Scalp and Neck⁴

ICD-10-CM CODE	DESCRIPTOR
C44.42	Squamous cell carcinoma of skin of scalp and neck

C44.52: Squamous Cell Carcinoma of Skin of Trunk⁴

ICD-10-CM CODE	DESCRIPTOR
C44.520	Squamous cell carcinoma of anal skin
C44.521	Squamous cell carcinoma of skin of breast
C44.529	Squamous cell carcinoma of skin of other part of trunk

Before prescribing KEYTRUDA, please read the Selected Safety Information on pages 5-8 and the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg

KEYTRUDA FOR RECURRENT OR METASTATIC CUTANEOUS SQUAMOUS CELL CARCINOMA (cSCC) (continued)

C44.62: Squamous Cell Carcinoma of Skin of Upper Limb, Including Shoulder⁴

ICD-10-CM CODE	DESCRIPTOR
C44.621	Squamous cell carcinoma of skin of unspecified upper limb, including shoulder
C44.622	Squamous cell carcinoma of skin of right upper limb, including shoulder
C44.629	Squamous cell carcinoma of skin of left upper limb, including shoulder

C44.72: Squamous Cell Carcinoma of Skin of Lower Limb, Including Hip⁴

ICD-10-CM CODE	DESCRIPTOR
C44.721	Squamous cell carcinoma of skin of unspecified lower limb, including hip
C44.722	Squamous cell carcinoma of skin of right lower limb, including hip
C44.729	Squamous cell carcinoma of skin of left lower limb, including hip

C44.8: Other and Unspecified Malignant Neoplasm of Overlapping Sites of Skin⁴

ICD-10-CM CODE	DESCRIPTOR
C44.82	Squamous cell carcinoma of overlapping sites of skin

C44.9: Other and Unspecified Malignant Neoplasm of Skin, Unspecified⁴

ICD-10-CM CODE	DESCRIPTOR
C44.92	Squamous cell carcinoma of skin, unspecified

CODING & BILLING

SELECTED SAFETY INFORMATION

Severe and Fatal Immune-Mediated Adverse Reactions

- KEYTRUDA is a monoclonal antibody that belongs to a class of drugs that bind to either the programmed death receptor-1 (PD-1) or the programmed death ligand 1 (PD-L1), blocking the PD-1/PD-L1 pathway, thereby removing inhibition of the immune response, potentially breaking peripheral tolerance and inducing immune-mediated adverse reactions. Immune-mediated adverse reactions, which may be severe or fatal, can occur in any organ system or tissue, can affect more than one body system simultaneously, and can occur at any time after starting treatment or after discontinuation of treatment. Important immune-mediated adverse reactions listed here may not include all possible severe and fatal immune-mediated adverse reactions.
- Monitor patients closely for symptoms and signs that may be clinical manifestations of underlying immune-mediated adverse reactions. Early identification and management are essential to ensure safe use of anti-PD-1/PD-L1 treatments. Evaluate liver enzymes, creatinine, and thyroid function at baseline and periodically during treatment. In cases of suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.
- Withhold or permanently discontinue KEYTRUDA depending on severity of the immune-mediated adverse reaction. In general, if KEYTRUDA requires interruption or discontinuation, administer systemic corticosteroid therapy (1 to 2 mg/kg/day prednisone or equivalent) until improvement to Grade 1 or less. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose adverse reactions are not controlled with corticosteroid therapy.

Immune-Mediated Pneumonitis

- KEYTRUDA can cause immune-mediated pneumonitis. The incidence is higher in patients who have received prior thoracic radiation. Immune-mediated pneumonitis occurred in 3.4% (94/2799) of patients receiving KEYTRUDA, including fatal (0.1%), Grade 4 (0.3%), Grade 3 (0.9%), and Grade 2 (1.3%) reactions. Systemic corticosteroids were required in 67% (63/94) of patients. Pneumonitis led to permanent discontinuation of KEYTRUDA in 1.3% (36) and withholding in 0.9% (26) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement; of these, 23% had recurrence. Pneumonitis resolved in 59% of the 94 patients.
- Pneumonitis occurred in 8% (31/389) of adult patients with cHL receiving KEYTRUDA as a single agent, including Grades 3-4 in 2.3% of patients. Patients received high-dose corticosteroids for a median duration of 10 days (range: 2 days to 53 months). Pneumonitis rates were similar in patients with and without prior thoracic radiation. Pneumonitis led to discontinuation of KEYTRUDA in 5.4% (21) of patients. Of the patients who developed pneumonitis, 42% interrupted KEYTRUDA, 68% discontinued KEYTRUDA, and 77% had resolution.

Immune-Mediated Colitis

- KEYTRUDA can cause immune-mediated colitis, which may present with diarrhea. Cytomegalovirus infection/reactivation has been reported in patients with corticosteroid-refractory immune-mediated colitis. In cases of corticosteroid-refractory colitis, consider repeating infectious workup to exclude alternative etiologies. Immune-mediated colitis occurred in 1.7% (48/2799) of patients receiving KEYTRUDA, including Grade 4 (<0.1%), Grade 3 (1.1%), and Grade 2 (0.4%) reactions. Systemic corticosteroids were required in 69% (33/48); additional immunosuppressant therapy was required in 4.2% of patients. Colitis led to permanent discontinuation of KEYTRUDA in 0.5% (15) and withholding in 0.5% (13) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement; of these, 23% had recurrence. Colitis resolved in 85% of the 48 patients.

Hepatotoxicity and Immune-Mediated Hepatitis

KEYTRUDA as a Single Agent

- KEYTRUDA can cause immune-mediated hepatitis. Immune-mediated hepatitis occurred in 0.7% (19/2799) of patients receiving KEYTRUDA, including Grade 4 (<0.1%), Grade 3 (0.4%), and Grade 2 (0.1%) reactions. Systemic corticosteroids were required in 68% (13/19) of patients; additional immunosuppressant therapy was required in 11% of patients. Hepatitis led to permanent discontinuation of KEYTRUDA in 0.2% (6) and withholding in 0.3% (9) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement; of these, none had recurrence. Hepatitis resolved in 79% of the 19 patients.

Before prescribing KEYTRUDA, please read the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg

CODING & BILLING

SELECTED SAFETY INFORMATION (*continued*)

Hepatotoxicity and Immune-Mediated Hepatitis (*continued*)

KEYTRUDA With Axitinib

- KEYTRUDA in combination with axitinib can cause hepatic toxicity. Monitor liver enzymes before initiation of and periodically throughout treatment. Consider monitoring more frequently as compared to when the drugs are administered as single agents. For elevated liver enzymes, interrupt KEYTRUDA and axitinib, and consider administering corticosteroids as needed. With the combination of KEYTRUDA and axitinib, Grades 3 and 4 increased alanine aminotransferase (ALT) (20%) and increased aspartate aminotransferase (AST) (13%) were seen at a higher frequency compared to KEYTRUDA alone. Fifty-nine percent of the patients with increased ALT received systemic corticosteroids. In patients with ALT ≥ 3 times upper limit of normal (ULN) (Grades 2-4, n=116), ALT resolved to Grades 0-1 in 94%. Among the 92 patients who were rechallenged with either KEYTRUDA (n=3) or axitinib (n=34) administered as a single agent or with both (n=55), recurrence of ALT ≥ 3 times ULN was observed in 1 patient receiving KEYTRUDA, 16 patients receiving axitinib, and 24 patients receiving both. All patients with a recurrence of ALT ≥ 3 ULN subsequently recovered from the event.

Immune-Mediated Endocrinopathies

Adrenal Insufficiency

- KEYTRUDA can cause primary or secondary adrenal insufficiency. For Grade 2 or higher, initiate symptomatic treatment, including hormone replacement as clinically indicated. Withhold KEYTRUDA depending on severity. Adrenal insufficiency occurred in 0.8% (22/2799) of patients receiving KEYTRUDA, including Grade 4 (<0.1%), Grade 3 (0.3%), and Grade 2 (0.3%) reactions. Systemic corticosteroids were required in 77% (17/22) of patients; of these, the majority remained on systemic corticosteroids. Adrenal insufficiency led to permanent discontinuation of KEYTRUDA in <0.1% (1) and withholding in 0.3% (8) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement.

Hypophysitis

- KEYTRUDA can cause immune-mediated hypophysitis. Hypophysitis can present with acute symptoms associated with mass effect such as headache, photophobia, or visual field defects. Hypophysitis can cause hypopituitarism. Initiate hormone replacement as indicated. Withhold or permanently discontinue KEYTRUDA depending on severity. Hypophysitis occurred in 0.6% (17/2799) of patients receiving KEYTRUDA, including Grade 4 (<0.1%), Grade 3 (0.3%), and Grade 2 (0.2%) reactions. Systemic corticosteroids were required in 94% (16/17) of patients; of these, the majority remained on systemic corticosteroids. Hypophysitis led to permanent discontinuation of KEYTRUDA in 0.1% (4) and withholding in 0.3% (7) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement.

Thyroid Disorders

- KEYTRUDA can cause immune-mediated thyroid disorders. Thyroiditis can present with or without endocrinopathy. Hypothyroidism can follow hyperthyroidism. Initiate hormone replacement for hypothyroidism or institute medical management of hyperthyroidism as clinically indicated. Withhold or permanently discontinue KEYTRUDA depending on severity. Thyroiditis occurred in 0.6% (16/2799) of patients receiving KEYTRUDA, including Grade 2 (0.3%). None discontinued, but KEYTRUDA was withheld in <0.1% (1) of patients.
- Hyperthyroidism occurred in 3.4% (96/2799) of patients receiving KEYTRUDA, including Grade 3 (0.1%) and Grade 2 (0.8%). It led to permanent discontinuation of KEYTRUDA in <0.1% (2) and withholding in 0.3% (7) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement. Hypothyroidism occurred in 8% (237/2799) of patients receiving KEYTRUDA, including Grade 3 (0.1%) and Grade 2 (6.2%). It led to permanent discontinuation of KEYTRUDA in <0.1% (1) and withholding in 0.5% (14) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement. The majority of patients with hypothyroidism required long-term thyroid hormone replacement. The incidence of new or worsening hypothyroidism was higher in 1185 patients with HNSCC, occurring in 16% of patients receiving KEYTRUDA as a single agent or in combination with platinum and FU, including Grade 3 (0.3%) hypothyroidism. The incidence of new or worsening hypothyroidism was higher in 389 adult patients with cHL (17%) receiving KEYTRUDA as a single agent, including Grade 1 (6.2%) and Grade 2 (10.8%) hypothyroidism.

Before prescribing KEYTRUDA, please read the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg

CODING & BILLING

SELECTED SAFETY INFORMATION (*continued*)

Immune-Mediated Endocrinopathies (*continued*)

Type 1 Diabetes Mellitus (DM), Which Can Present With Diabetic Ketoacidosis

- Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold KEYTRUDA depending on severity. Type 1 DM occurred in 0.2% (6/2799) of patients receiving KEYTRUDA. It led to permanent discontinuation in <0.1% (1) and withholding of KEYTRUDA in <0.1% (1) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement.

Immune-Mediated Nephritis With Renal Dysfunction

- KEYTRUDA can cause immune-mediated nephritis. Immune-mediated nephritis occurred in 0.3% (9/2799) of patients receiving KEYTRUDA, including Grade 4 (<0.1%), Grade 3 (0.1%), and Grade 2 (0.1%) reactions. Systemic corticosteroids were required in 89% (8/9) of patients. Nephritis led to permanent discontinuation of KEYTRUDA in 0.1% (3) and withholding in 0.1% (3) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement; of these, none had recurrence. Nephritis resolved in 56% of the 9 patients.

Immune-Mediated Dermatologic Adverse Reactions

- KEYTRUDA can cause immune-mediated rash or dermatitis. Exfoliative dermatitis, including Stevens-Johnson syndrome, drug rash with eosinophilia and systemic symptoms, and toxic epidermal necrolysis, has occurred with anti-PD-1/PD-L1 treatments. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate nonexfoliative rashes. Withhold or permanently discontinue KEYTRUDA depending on severity. Immune-mediated dermatologic adverse reactions occurred in 1.4% (38/2799) of patients receiving KEYTRUDA, including Grade 3 (1%) and Grade 2 (0.1%) reactions. Systemic corticosteroids were required in 40% (15/38) of patients. These reactions led to permanent discontinuation in 0.1% (2) and withholding of KEYTRUDA in 0.6% (16) of patients. All patients who were withheld reinitiated KEYTRUDA after symptom improvement; of these, 6% had recurrence. The reactions resolved in 79% of the 38 patients.

Other Immune-Mediated Adverse Reactions

- The following clinically significant immune-mediated adverse reactions occurred at an incidence of <1% (unless otherwise noted) in patients who received KEYTRUDA or were reported with the use of other anti-PD-1/PD-L1 treatments. Severe or fatal cases have been reported for some of these adverse reactions. *Cardiac/Vascular*: Myocarditis, pericarditis, vasculitis; *Nervous System*: Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis (including exacerbation), Guillain-Barré syndrome, nerve paresis, autoimmune neuropathy; *Ocular*: Uveitis, iritis and other ocular inflammatory toxicities can occur. Some cases can be associated with retinal detachment. Various grades of visual impairment, including blindness, can occur. If uveitis occurs in combination with other immune-mediated adverse reactions, consider a Vogt-Koyanagi-Harada-like syndrome, as this may require treatment with systemic steroids to reduce the risk of permanent vision loss; *Gastrointestinal*: Pancreatitis, to include increases in serum amylase and lipase levels, gastritis, duodenitis; *Musculoskeletal and Connective Tissue*: Myositis/polymyositis, rhabdomyolysis (and associated sequelae, including renal failure), arthritis (1.5%), polymyalgia rheumatica; *Endocrine*: Hypoparathyroidism; *Hematologic/Immune*: Hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis, systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenic purpura, solid organ transplant rejection.

Infusion-Related Reactions

- KEYTRUDA can cause severe or life-threatening infusion-related reactions, including hypersensitivity and anaphylaxis, which have been reported in 0.2% of 2799 patients receiving KEYTRUDA. Monitor for signs and symptoms of infusion-related reactions. Interrupt or slow the rate of infusion for Grade 1 or Grade 2 reactions. For Grade 3 or Grade 4 reactions, stop infusion and permanently discontinue KEYTRUDA.

Before prescribing KEYTRUDA, please read the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg

CODING & BILLING

SELECTED SAFETY INFORMATION (*continued*)

Complications of Allogeneic Hematopoietic Stem Cell Transplantation (HSCT)

- Fatal and other serious complications can occur in patients who receive allogeneic HSCT before or after anti-PD-1/PD-L1 treatments. Transplant-related complications include hyperacute graft-versus-host disease (GVHD), acute and chronic GVHD, hepatic veno-occlusive disease after reduced intensity conditioning, and steroid-requiring febrile syndrome (without an identified infectious cause). These complications may occur despite intervening therapy between anti-PD-1/PD-L1 treatments and allogeneic HSCT. Follow patients closely for evidence of these complications and intervene promptly. Consider the benefit vs risks of using anti-PD-1/PD-L1 treatments prior to or after an allogeneic HSCT.

Increased Mortality in Patients With Multiple Myeloma

- In trials in patients with multiple myeloma, the addition of KEYTRUDA to a thalidomide analogue plus dexamethasone resulted in increased mortality. Treatment of these patients with an anti-PD-1/PD-L1 treatment in this combination is not recommended outside of controlled trials.

Embryofetal Toxicity

- Based on its mechanism of action, KEYTRUDA can cause fetal harm when administered to a pregnant woman. Advise women of this potential risk. In females of reproductive potential, verify pregnancy status prior to initiating KEYTRUDA and advise them to use effective contraception during treatment and for 4 months after the last dose.

Adverse Reactions

- When KEYTRUDA was used as monotherapy, the most common adverse reactions ($\geq 20\%$) were fatigue, musculoskeletal pain, decreased appetite, pruritus, diarrhea, nausea, rash, pyrexia, cough, dyspnea, constipation, pain, and abdominal pain.
- When KEYTRUDA was used in combination with chemotherapy, the most common adverse reactions ($\geq 20\%$) were fatigue/asthenia, nausea, constipation, diarrhea, decreased appetite, rash, vomiting, cough, dyspnea, pyrexia, alopecia, peripheral neuropathy, mucosal inflammation, stomatitis, headache, and weight loss.
- When KEYTRUDA was used in combination with axitinib, the most common adverse reactions ($\geq 20\%$) were diarrhea, fatigue/asthenia, hypertension, hepatotoxicity, hypothyroidism, decreased appetite, palmar-plantar erythrodysesthesia, nausea, stomatitis/mucosal inflammation, dysphonia, rash, cough, and constipation.
- When KEYTRUDA was used in combination with LENVIMA, the most common adverse reactions ($\geq 20\%$) were fatigue, musculoskeletal pain, hypertension, diarrhea, decreased appetite, hypothyroidism, nausea, stomatitis, vomiting, weight loss, abdominal pain, headache, constipation, urinary tract infection, dysphonia, hemorrhagic events, hypomagnesemia, palmar-plantar erythrodysesthesia syndrome, dyspnea, cough, and rash.

Lactation

- Because of the potential for serious adverse reactions in breastfed children, advise women not to breastfeed during treatment and for 4 months after the final dose.

Pediatric Use

- In KEYNOTE-051, 161 pediatric patients (62 pediatric patients aged 6 months to younger than 12 years and 99 pediatric patients aged 12 years to 17 years) were administered KEYTRUDA 2 mg/kg every 3 weeks. The median duration of exposure was 2.1 months (range: 1 day to 24 months).
- Adverse reactions that occurred at a $\geq 10\%$ higher rate in pediatric patients when compared to adults were pyrexia (33%), vomiting (30%), leukopenia (30%), upper respiratory tract infection (29%), neutropenia (26%), headache (25%), and Grade 3 anemia (17%).

cHL=classical Hodgkin lymphoma; FU=fluorouracil; HNSCC=head and neck squamous cell carcinoma.

Before prescribing KEYTRUDA, please read the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg